



Random walks: the drunk and the canyon - Tutorial for R



Imagine that a drunkard is walking in a huge plains, that has a canyon on one of the sides. Every time the drunkard steps forward, he also staggers to the left or to the right, with equal probability.

This is a very simple Markovian process, called [random walk](#), on one dimension¹⁾. When the drunkard falls in the canyon, the random walk is over (and so is the drunkard), so we call this condition as *absorbing boundary*.

Virtual drunks

To proceed, you must have the R environment with the Ecovirtual package installed and loaded. If you do not have and do not know how to have them, see the [Installation page](#).

What can we predict about this process? Let's set some drunks lose on this virtual plains. To do this, we will use the function **randWalk** from EcoVirtual R package.

Simulation parameters:

Option	Parameter	
Number of Species	S	Number of drunks
Step Size	step	Number of steps the drunk moves sideways at each time interval

Option	Parameter	
Maximum Initial Distance	xlmax	maximum distance between the drunks and the canyon when the simulation starts
Initial Distance Equal	alleg=TRUE	when selected (TRUE), all drunks start at the maximum distance. When unselected, the drunks start at random position from 1 to the maximum distance above
Maximum time	tmax	total simulation time

Example

Let's simulate ten drunkards, staggering for 10 steps at each time interval, for ten thousand time steps:

```
S = 10
step = 10
xlmax = 200
alleg = TRUE
tmax = 10000
```

As with every stochastic process, the results may vary at each simulation. Thus, repeat the simulation some times to make sure you understand the results.

Step effect

What happens if the drunks are more or less staggering? Try reducing the step size to 2:

```
S = 10
step = 2
xlmax = 200
alleg = TRUE
tmax = 10000
```

Time effect

Drunks that stagger less are less prone to end in the bottom of the canyon, or is it only a matter of time? Try increasing the final time:

```
S = 10
step = 2
xlmax = 200
alleg = TRUE
tmax = 50000
```

Question

The drunk has equal probability to stagger left or right, so on the average he only walks forward.

Given enough time, this random walk with absorbing boundary has only one possible outcome. What is it?

Virtual population

\ To simulate this dynamics of birth and death, use the function **estDem** from EcoVirtual R package.

The options here control the simulation of populations under a continuous time random walk:

Option	Parameter	Definition
Enter name for last simulation data set	R object	R object in which to save the simulation results
Maximum time	tmax	Maximum simulation time, in the same time unit as the rates used
Number of simulations	nsim	Number of different populations to simulate
Initial size	N0	Initial population size
birth rate	b	instantaneous birth rate (\$b\$)
death rate	d	instantaneous death rate (\$d\$)

Example

Simulate the trajectory of 20 population with equal birth and death rates, starting with 10 individuals each. Let the time continue up to 50 units:

```
tmax = 50
nsim = 20
N0 = 10
b = 0.2
d = 0.2
```

You should see a random walk graph, very similar to the drunks. The number of extinctions up to the Maximum time is recorded in the upper left corner of the graph.

Questions

1. Which are the corresponding parameters between the drunk random walk and the birth and death dynamics?
2. The effects you have noted on the drunk simulation (step size and end time) affect the population dynamics in the same way?
3. What is the consequence of this result for the conservation of natural populations?

To learn more

- We have simulated a dynamic with equal probability of births and deaths with an absorbing boundary. This is a particular case of stochastic birth and death processes. You can read more

about them in the [tutorial on demographic stochasticity](#).

- [Chemotaxis - How a Small Organism Finds a Food Source](#): one excellent explanation about random walk and its application to another area in biology, as presented by MIT students.

1)

we are only interested in the lateral movement

From:

<http://ecovirtual.ib.usp.br/> -

Permanent link:

<http://ecovirtual.ib.usp.br/doku.php?id=en:ecovirt:roteiro:math:bebador>



Last update: **2017/09/19 17:17**