

- 
- 
- 

Metapopulações com colonização interna - Roteiro no EcoVirtual



There's an assumption in the last model ([Metapopulations and propagule rain - Tutorial in EcoVirtual](#)) that we can get rid off: the assumption that the propagule rain is constant. Instead, we can make the colonization rate a function of the percentage of occupied patches. In a simple formulation of this model, the propagule source is just from within the system (closed system) and the probability of colonization increases linearly with the proportion of occupied patches.

The model

This way, our model does not have a constant colonization rate p_i , but a probability of colonization that depends on the number of occupied patches:

$$p_i = \text{if} \{$$

here i is a constant that indicates how much the colonization probability increases with every new occupied patch. This way, the more occupied patches we have, the greater is the probability that new patches will be occupied. Substituting this new formula for p_i in our old equation, we have:

$$\frac{df}{dt} = \text{if}(1-f) - p_e f$$

Equilibrium

The fraction of the occupied patches in equilibrium (F) changes to:


$$F = \frac{p_e}{1 + p_e}$$

Simulação

Let's see this first-hand by simulating this situation. As in the last exercise, we have created an R function that runs the simulation for us. As before, this function simply draws random colonization and extinction events for every patch at every time step, following the model's rules. It then shows the graph and an occupancy matrix for every time interval.

To proceed, you must have the R environment with the Rcmdr and Ecovirtual packages installed and loaded. If you do not have and do not know how to have them, see the [Installation page](#).

To run this model in **EcoVirtual** use the function ***Internal colonization*** do sub-menu **Metapopulation** in the Rcommander/EcoVirtual menu.



The parameters are:

option	parameter	definition
data set	R object	where the results are stored
Maximum time	\$tmax\$	Number of model iterations
columns	\$cl\$	number of habitat columns in the landscape

option	parameter	definition
rows	\$rw\$	number of habitat rows in the landscape
initial occupancy	\$f0\$	initial number of occupied patches
colonization coef.	\$ci\$	colonization coefficient
prob. extinction	\$pe\$	extinction probability

Now you can simulate the model with the values you chose. For example,

```
tmax = 100
ncol = 10
nrow = 10
f0 = 0.1
ci = 1
pe = 0.5
```

Play around with the model for a while by changing the parameters and try to answer the following questions:

- is there any difference between this model and the *seed rain*, keeping the shared parameters equal?
- does the position of a patch in the landscape affect its \$p_i\$ or \$p_e\$? Is this realistic?
- are there any combinations of \$i\$ and \$p_e\$ that cannot exist?
- what is the meaning of a negative F ?
- in which situation the equilibrium is $\hat{f} = 1$?

Scenario suggestion

```
tmax = 100
cl = 10
rw = 10
f0 = 0.1
ci = 0.5
pe = 0.5
```

To learn more

- **Gotelli, N. 2007. Ecologia.** Londrina, Ed. Planta. Capítulo 4.
- **Stevens, M. H. 2009. A primer of ecology with R.** New York. Springer. Capítulo 4.
- **Gotelli, N. 1991. Metapopulation models: the rescue effect, the propagule rain, and the core-satellite hypothesis.** The American Naturalist, 138: 768-776. [pdf no site do autor](#)

[RCMDR, uma população, metapopulações, colonização interna](#)

Last
update:
2017/10/10 en:ecovirt:roteiro:metap_uma:metap_circmdr http://ecovirtual.ib.usp.br/doku.php?id=en:ecovirt:roteiro:metap_uma:metap_circmdr&rev=1507670016
18:13

From:
<http://ecovirtual.ib.usp.br/> -

Permanent link:
http://ecovirtual.ib.usp.br/doku.php?id=en:ecovirt:roteiro:metap_uma:metap_circmdr&rev=1507670016

Last update: **2017/10/10 18:13**