




Successional Niche - Tutorial for Ecovirtual



Usually the competition between species is viewed as a “all or nothing” instantaneous process. However, observing nature we realize that plants with high capacity for colonization generally have high metabolic rates (respiration, photosynthesis and allocation of reproductive tissue). These high rates allow for the plants to grow and reproduce more quickly, which may give them an extra advantage in competitive interaction. Imagine a forest where a clearing was opened by a fallen tree and both species, the best competitor and the better disperser reach it at the same time. In this situation, imagining that the best competitor will immediately remove the other does not seem very reasonable, simply because there is no resource limitation yet. On the other hand, the species that have the highest growth rate can reproduce before the resource limitation occurs and it is driven to local extinction by competition.

Pacala and Rees model


This period, before the environmental resource reduction, creates an ephemeral niche that was called by Pacala and Rees (1998) the “succession niche”. These authors developed a simple model to test their ideas. They have established five possible states in the system:

1. Vacant: none of the species
2. Early: occupied only by the species of early succession
3. Susceptible: occupied by the late species, but susceptible to initial invasion because the resources are still abundant
4. Mixed: occupied by both species, on the road to competitive exclusion
5. Resistant: occupied only by the late species and resistant to invasion

Given these states the succession process can have several possible trajectories:

- VACANT → EARLY → MIXED → RESISTANT
- VACANT → SUSCEPTIBLE → MIXED → RESISTANT
- VACANT → SUSCEPTIBLE → RESISTANT

Let's create our model. For simplicity, rather than we model each species, we model the state and its transitions in a similar way that we model the states of individuals in a population: remember the matrix models of Leslie and Lefkovich from the earlier classes?! Look at the diagram below to understand the state transitions:



This model has four parameters: c , α , m and γ :

- c is the base colonization rate
- α is the early species relative colonization rate
- m is the death or disturbance rate
- γ is the rate of competitive exclusion


With these four parameters, we can model the variation of the proportion of states over time, with the expressions that appear in the transitions from the figure above. Solid lines indicate increase in proportion and broken lines decreases in proportion. For example, variation in the SUSCEPTIBLE state is given by:

$$\frac{dS}{dt} = [c(S + R + M)]V - [\alpha c(M+E)]S - gS - mS$$

Using o EcoVirtual

To proceed, you must have the R environment with the Rcmdr and Ecovirtual packages installed and loaded. If you do not have and do not know how to have them, see the [Installation page](#).

Let's use EcoVirtual to simulate this model's dynamics. Open the menu **EcoVirtual>Multi species>Niche regeneration...**:



The following parameters can be changed:

option	parameter	effect
data set	R object	stores the simulation results
Simulation Arena Condition	Basic simulation parameters	
Maximum time	tmax	number of model iterations to be run
columns	cl	number of columns in the habitat
rows	rw	number of rows in the habitat
Initial Stages Proportions	Initial state	
Early Stage	er	fraction of patches initially occupied by <i>sp2</i>
Susceptible	/sc	fraction of patches initially with <i>sp1</i> , and that may be invaded by <i>sp2</i>
Mixed	mx	fraction of patches initially with both species
Resistant	rs	fraction of patches initially with <i>sp1</i> , and that may not be colonized <i>sp2</i>
Colonization rates	Colonization parameters	
Better competitor	c1	<i>sp1</i> colonization coefficient
Poor competitor	c2	<i>sp2</i> colonization coefficient
General Parameters	General parameters	
Competitive exclusion	ec	probability of transition from SC and MX to RS
Disturbance	dst	proportion of patches that is kept empty

Testing with a high rate of competitive exclusion and low disturbance:

```
tmax=50,  
rw=100,  
cl=100,  
c1=0.2,  
c2=0.8,  
ec=0.5,  
dst=0.04,  
er=0.08,  
sc=0.02,  
mx=0,  
rs=0,
```

Now, let's simulate some scenarios to see what happens:

1. Reduce the competitive exclusion to 0.1.
2. Keep *ec* in 0.1 and increase the disturbance to 0.1 as well.
3. Put both species with the same colonization rate (*c1*=0.4 and *c2*=0.4).

What is the biological interpretation for your results? Connect your answer to:

- ecological succession
- successional niche
- intermediate disturbance
- competition

Questions

1. Write a short essay in which you provide an interpretation for all of the scenarios above, and connect the trajectories to the concepts of:
 - ecological succession
 - successional niche
 - intermediate disturbance
 - competition
1. How can you change the function used to create a scenario in which you would only study the tradeoff between colonization vs. competition? Explain your answer.
2. Is coexistence possible if you only consider the effect of the successional niche? Use a simulation to justify your answer.

To learn more

- [Pacala, S. & M. Rees. 1998. Models suggesting field experiments to test two hypotheses explaining successional diversity. The American Naturalist 152\(2\): 729:737.](#)
- [Stevens, M.H.H. \(2009\) A primer in ecology with R. New York, Springer. capítulo 9](#)

[RCMDR](#), [comunidades](#), [sucessão](#)

From:

<http://ecovirtual.ib.usp.br/> -

Permanent link:

http://ecovirtual.ib.usp.br/doku.php?id=en:ecovirt:roteiro:sucess:nich_regrcmdr



Last update: **2017/11/01 10:36**